

Abstract Submitted  
for the DAMOP17 Meeting of  
The American Physical Society

**Control of photoemission delay in resonant two-photon transitions**<sup>1</sup>

L. ARGENTI, Univ. Central Florida, FL, USA; Univ. Autonoma de Madrid, ES, EU (UAM), A. JIMENEZ GALAN, Max Born Inst., Berlin, D, EU; UAM, R. TAIEB, J. CAILLAT, A. MAQUET, Sorbonne Univ.; UPMC Univ. Paris 6, FR, EU, F. MARTIN, UAM; IMDEA-Nanociencia, IFIMAC, Madrid, ES, EU — The emission time delay  $\tau$  in one-photon absorption, which coincides with half the Wigner scattering delay  $\tau_W$ , is a fundamental descriptor of the photoelectric effect. While it is hard to access  $\tau$  in a direct way, it is possible to extrapolate it from the delay in two-photon transitions,  $\tau^{(2)}$ , measured with attosecond pump-probe schemes, provided that the contribution of the probe stage can be factored out. In absence of resonances,  $\tau$  can be expressed as the energy derivative of the dipole ionization amplitude,  $\tau = \partial_E \arg D_{Eg}$ , and  $\tau \simeq \tau^{(2)} - \tau_{cc}$  where  $\tau_{cc}$  is associated to the dipole transition in the continuum. Here we show that in the presence of a resonance the correspondence between  $\tau$  and  $\partial_E \arg D_{Eg}$  is lost. Furthermore, while  $\tau^{(2)}$  still coincides with  $\partial_E \arg D_{Eg}^{(2)}$ , it does not have any scattering counterpart. Indeed,  $\tau^{(2)}$  can be much larger than the lifetime of an intermediate resonance in the two-photon process, or more negative than the lower bound imposed on scattering delays by causality. Finally,  $\tau^{(2)}$  is controlled by the probe frequency. By varying  $\omega_{IR}$ , therefore, it is possible to radically alter a photoelectron group delay.

<sup>1</sup>NSF Grant No. 1607588, ERC Grant No. 290853 XCHEM, MINECO FIS2013-42002-R/FIS2016-77889-R, e-COST Action XLIC CM1204.

Luca Argenti  
Univ of Central Florida

Date submitted: 26 Jan 2017

Electronic form version 1.4