

Presenting both side-by-side on Wed or one on Wed and one on Thurs.  
would be  
optimal.

Abstract Submitted  
for the DAMOP17 Meeting of  
The American Physical Society

**Theoretical Studies of Dissociative Recombination of Electrons with SH<sup>+</sup> Ions**<sup>1</sup> D. O. KASHINSKI<sup>2</sup>, O. E. DI NALLO, United States Military Academy, A. P. HICKMAN, Lehigh University, J. ZS. MEZEI, F. COLBOC, I. F. SCHNEIDER, Université du Havre, K. CHAKRABARTI, University of Kolkatta, D. TALBI, Université Montpellier — We are investigating the dissociative recombination (DR) of electrons with the molecular ion SH<sup>+</sup>, i.e.  $e^- + \text{SH}^+ \rightarrow \text{S} + \text{H}$ . SH<sup>+</sup> is found in the interstellar medium (ISM), and little is known concerning its chemistry. Understanding the role of DR of electrons with SH<sup>+</sup> will lead to more accurate astrophysical models. Large active-space multi-reference configuration interaction (MRCI) electronic structure calculations were performed using the GAMESS code to obtain ground and excited <sup>2</sup>Π state potential energy curves (PECs) for several values of SH separation. Core-excited Rydberg states have proven to be of huge importance. The block diagonalization method was used to disentangle interacting states and form a diabatic representation of the PECs. Currently we are performing dynamics calculations using Multichannel Quantum Defect Theory (MQDT) to obtain DR rates. The status of the work will be presented at the conference.

<sup>1</sup>work supported by the French CNRS, the NSF, the XSEDE, and USMA

<sup>2</sup>travel supported by DoD-HPCMP

D. O. Kashinski  
United States Military Academy

Date submitted: 27 Jan 2017

Electronic form version 1.4