

Abstract Submitted  
for the DAMOP17 Meeting of  
The American Physical Society

**Direct search for the thorium-229 nuclear isomeric transition with a pulsed VUV laser** JUSTIN JEET, CHRISTIAN SCHNEIDER, University of California, Los Angeles, EUGENE V. TKALYA, Lomonosov Moscow State University and Nuclear Safety Institute of Russian Academy of Science, ERIC R. HUDSON, University of California, Los Angeles — The nucleus of thorium-229 has an exceptionally low-energy isomeric transition in the vacuum-ultraviolet (VUV) spectrum around  $7.8 \pm 0.5\text{eV}$  [1]. While inaccessible to standard nuclear physics techniques, there are various prospects for a laser-accessible nuclear transition. Our direct search for the transition uses thorium-doped crystals as samples. In a previous experiment [2] at the Advanced Light Source (ALS) synchrotron, LBNL, we were able to exclude a large portion of the transition lifetime-vs.-frequency region-of-interest (ROF) [3]. Here, we will report on our ongoing efforts of a search using a pulsed VUV laser system as light source, which allows us to enhance our sensitivity up to  $10^4\times$  over the ALS and extend the accessible frequency range over the entire ROF [3]. An updated exclusion region will be presented.

- [1] B. R. Beck et al.: LLNL-PROC-415170 (2009)
- [2] J. Jeet et al.: Phys. Rev. Lett. 114, 253001 (2015)
- [3] E. V. Tkalya et al.: Phys. Rev. C 92, 054324 (2015)

Christian Schneider  
University of California, Los Angeles

Date submitted: 28 Jan 2017

Electronic form version 1.4