

Abstract Submitted
for the DAMOP17 Meeting of
The American Physical Society

Vortex cluster shedding in an oblate Bose-Einstein condensate¹

YOUNGHOON LIM, JUNHONG GOO, WOO JIN KWON, JOON HYUN KIM, SANG WON SEO, YONG-IL SHIN, Center for Correlated Electron Systems, Institute for Basic Science (IBS), Seoul 08826, Republic of Korea — We present the observation of vortex cluster shedding from a moving obstacle in an oblate Bose-Einstein condensate (BEC). We investigate the evolution of the vortex shedding pattern as a function of the obstacle velocity v , and observe regular shedding of vortex clusters consisting of two like-sign vortices at low obstacle velocity. As v increases, the vortex shedding pattern becomes irregular with many larger clusters, which shows a transition to turbulence. To quantitatively characterize the vortex shedding pattern, we analyze the cluster charge distribution as a function of the obstacle velocity. The transition from regular to turbulent shedding is manifested with a rapid decrease of fractional population of two like-sign vortex clusters. In particular, we observe a saturating behavior of the Strouhal number with increasing v , which is associated with the shedding frequency. We will discuss possible extension of this work to test the universality of the vortex shedding dynamics.

¹This work was supported by IBS-R009-D1

Younghoon Lim
Seoul Natl Univ

Date submitted: 09 Mar 2017

Electronic form version 1.4