

Abstract Submitted
for the DAMOP18 Meeting of
The American Physical Society

Towards quantum many-body physics with Sr in optical lattices

SEBASTIAN BLATT, ANDR HEINZ, ANNIE JIHYUN PARK, FABIAN FINGER, STEPAN SNIGIREV, Max-Planck-Institut fr Quantenoptik, Hans-Kopfermann-Strae 1, 85784 Garching, Germany, JEAN DALIBARD, Laboratoire Kastler Brossel, Collge de France, ENS-PSL Research University, CNRS, UPMC-Sorbonne Universits, 11 place Marcelin Berthelot, 75005 Paris, IMMANUEL BLOCH, Max-Planck-Institut fr Quantenoptik, Hans-Kopfermann-Strae 1, 85784 Garching, Germany — The widespread use of ultracold fermionic strontium atoms in optical lattices for precision measurements has led to the availability of many advanced tools and techniques for these atoms. With the recent realization of degenerate gases of all Sr isotopes and the development of fermionic quantum gas microscopes for alkali metal atoms, a new frontier has opened for quantum simulations and quantum information processing with fermionic ^{87}Sr . Many applications in quantum state engineering and quantum simulation require internal-state-dependent control of the atomic motion. In the Sr atom, there exist so-called tuneout wavelengths, where only one of the clock states is trapped and the other state can move freely. Because of the (in principle) exact cancellation of the other state's polarizability at the tuneout wavelengths, it should be possible to realize spin-dependent lattices with high fidelity. Here, we propose a system to realize such internal-state-dependent control of the atomic motion and report on the construction of a new experiment towards quantum simulations with Sr in optical lattices.

Sebastian Blatt
Max-Planck-Institut fr Quantenoptik

Date submitted: 24 Jan 2018

Electronic form version 1.4