

Abstract Submitted
for the DAMOP18 Meeting of
The American Physical Society

Excited state positronium formation from C_{60} and C_{240} molecules¹

PAUL-ANTOINE HERVIEUX, Universite de Strasbourg, CNRS, Institut de Physique et Chimie des Materiaux de Strasbourg, France, ANZUMAAN CHAKRABORTY, HIMADRI CHAKRABORTY, Northwest Missouri State University, Maryville, USA — There exists a broad landscape of positronium (Ps) formation studies by implanting positron beams on atoms, molecules, polymers, solids, liquids, surfaces/films, metal-organic-frameworks and embedded nanostructures. Recently, ground state Ps formation studies using the C_{60} molecule as a target have been carried out [1,2]. Due to the dominant electron capture from the fullerene shell region and the spatial dephasing across the shell-width, a powerful diffraction effect was found to underpin the process of Ps formation in C_{60} . In the current work, we access excited state Ps formations from both C_{60} and C_{240} fullerenes. Results reveal significant structures in the cross section as a function of both the impact energy and the Ps level – structures that arise from positron-fullerene short-range interactions. Fullerenes are modeled by a local-density approximation (LDA) approach [3] and the Ps formation is treated by the continuum distorted-wave final-state (CDW-FS) approximation [4]. The work may motivate application of the Ps formation spectroscopy to gas phase nanoparticles and access fullerene-level- as well as Ps-level-differential measurements. [1] Hervieux et al., Phys. Rev. A **95**, 020701 (R) (2017); [2] Chakraborty et al., J. Phys.: Conf. Series **875**, 072002 (2017); [3] Choi et al., Phys. Rev. A **95**, 023404 (2017); [4] Fojon et al., Phys. Rev. A **54**, 4923 (1996).

¹The work is supported by the US National Science Foundation.

Himadri Chakraborty
Northwest Missouri State University

Date submitted: 24 Jan 2018

Electronic form version 1.4