Abstract Submitted for the DFD07 Meeting of The American Physical Society

Anisotropic low-wavenumber constraints on energy in rotating and stratified flows<sup>1</sup> SUSAN KURIEN, BETH WINGATE, Los Alamos National Laboratory, MARK TAYLOR, Sandia National Laboratories — Rapidly rotating, stably stratified three-dimensional inviscid flows conserve both energy and potential enstrophy. We show that in such flows, the forward cascade of potential enstrophy imposes anisotropic constraints on the wavenumber distribution of kinetic and potential energy. The horizontal kinetic energy is suppressed in the large, nearly horizontal wave modes, and should decay with the horizontal wavenumber as  $k_h^{-3}$ . The potential energy is suppressed in the large, nearly vertical wave modes, and should decay with the vertical wavenumber as  $k_z^{-3}$  . These results augment the only other exact prediction for the scaling of energy spectra due to constraints by potential enstrophy obtained by Charney (J. Atmos. Sci. 28, 1087 (1971)), who showed that in the quasi-geostrophic approximation for rotating stratified flows, the energy spectra must scale isotropically with total wavenumber as  $k^{-3}$ . We test our predicted scaling estimates using resolved numerical simulations of the Boussinesq equations in the relevant parameter regimes, and find reasonable agreement.

<sup>1</sup>Supported by DOE Office of Science Applied Mathematics Program

Susan Kurien Los Alamos National Laboratory

Date submitted: 31 Jul 2007

Electronic form version 1.4