

Abstract Submitted
for the DFD11 Meeting of
The American Physical Society

Heat transport near the transition to the ultimate state of turbulent Rayleigh-Bénard convection¹ GUENTER AHLERS, UC Santa Barbara, XIAOZHOU HE, MPIDS Göttingen, Germany, DENIS FUNFSCHILLING, LSGC CNRS Nancy, France, HOLGER NOBACH, EBERHARD BODENSCHATZ, MPIDS Göttingen, Germany — Measurements of the Nusselt number Nu for Rayleigh-Bénard convection (RBC) of a cylindrical sample over the Rayleigh-number range $2 \times 10^{12} \leq Ra \leq 2 \times 10^{15}$ and the Prandtl-number range $0.79 \leq Pr \leq 0.86$ are presented. The aspect ratio $\Gamma \equiv D/L$ was 0.50 ($D = 1.12$ m was the diameter and $L = 2.24$ m was the height). For $Ra \leq 2 \times 10^{13}$ the data yielded $Nu = Nu_0 Ra^{\gamma_{eff}}$ with $\gamma_{eff} = 0.312$, consistent with classical RBC.² For larger Ra γ_{eff} increased, reaching approximately $\gamma_{eff} = 0.38$ near $Ra = 10^{15}$, consistent with predictions^{3,4} for an ultimate state with turbulent boundary layers.

¹Supported by the Max Planck Society, the Volkswagenstiftung, and NSF Grant DMR07-02111.

²G. Ahlers, S. Grossmann, and D. Lohse, *Rev. Mod. Phys.* **81**, 503 (2009)

³R. H. Kraichnan, *Phys. Fluids* **5**, 1374 (1962)

⁴S. Grossmann and D. Lohse, *Phys. Fluids* **23**, 045108 (2011)

Guenter Ahlers
UCSB

Date submitted: 05 Aug 2011

Electronic form version 1.4