

Abstract Submitted
for the DFD12 Meeting of
The American Physical Society

Soret-driven convection in colloidal suspensions MAHMOUD

DARASSI, LAYACHI HADJI, The University of Alabama — Convection in colloidal suspensions of solid particles is characterized by the interplay between thermophoresis, sedimentation and Brownian diffusion. Their coupled effects is represented by a dimensionless parameter β and experiments by Chang *et al.* (2008) have shown that for a given set of experimental parameters, β is a function of the particle radius r_p with the function $\beta(r_p)$ having the shape of an inverted parabola with two roots in the range $5 \text{ nm} \leq r_p \leq 125 \text{ nm}$. We investigate both the linear and non-linear convection in a suspension of solid particles using a particulate medium in a Rayleigh-Bénard geometry set-up. The analysis focuses on the particle dominated convection regime for which the onset is steady and to disturbances having infinitely long wavelength. For $0 < \beta \ll 1$, which corresponds to particle size near the two roots of $\beta(r)$, we retrieve the instability threshold conditions for the binary mixture model. For $\beta = O(1)$, we show that, unlike the binary mixture model, the conditions for instability onset can be mapped to corresponding experimental parameters. A nonlinear evolution equation is derived and its predictions compared to those of a similar equation for the binary mixture case.

Layachi Hadji
The University of Alabama

Date submitted: 01 Aug 2012

Electronic form version 1.4