

Abstract Submitted  
for the DFD15 Meeting of  
The American Physical Society

**Imbibition of “Open Capillary”: Fundamentals and Applications<sup>1</sup>**

MARIE TANI, Department of Physics, Faculty of Science, Ochanomizu University, RYUJI KAWANO, Division of Biotechnology and Life Science, Tokyo University of Agriculture and Technology, KOKI KAMIYA, Artificial Cell Membrane Systems Group, Kanagawa Academy of Science and Technology (KAST), KO OKUMURA, Department of Physics, Faculty of Science, Ochanomizu University — Control or transportation of small amount of liquid is one of the most important issues in various contexts including medical sciences or pharmaceutical industries to fuel delivery. We studied imbibition of “open capillary” both experimentally and theoretically, and found simple scaling laws for both statics and dynamics of the imbibition, similarly as that of imbibition of capillary tubes. Furthermore, we revealed the existence of “precursor film,” which developed ahead of the imbibing front, and the dynamics of it is described well by another scaling law for capillary rise in a corner [1]. Then, to show capabilities of open capillaries, we demonstrated two experiments by fabricating micro mixing devices to achieve (1) simultaneous multi-color change of the Bromothymol blue (BTB) solution and (2) expression of the green fluorescent protein (GFP) [2]. [1] A. Ponomarenko, D. Quere and C. Clanet, *J. Fluid Mech.*, 666, 146 (2011). [2] M. Tani, R. Kawano, K. Kamiya and K. Okumura, *Sci. Rep.* 5, 10263 (2015).

<sup>1</sup>This research was partly supported by ImPACT Program of Council for Science, Technology and Innovation (Cabinet Office, Government of Japan). M. T. is supported by the Japan Society for the Promotion of Science Research Fellowships for Young Scientists.

Marie Tani  
Department of Physics, Faculty of Science, Ochanomizu University

Date submitted: 16 Jul 2015

Electronic form version 1.4