

Abstract Submitted
for the DFD15 Meeting of
The American Physical Society

Right Heart Vorticity and Right Ventricular Diastolic Dysfunction JAMES BROWNING, JEAN HERTZBERG, University of Colorado Boulder, BRETT FENSTER, National Jewish Health and University of Colorado Denver, JOYCE SCHROEDER, University of Colorado Denver — Recent advances in cardiac magnetic resonance imaging (CMR) have allowed for the 3-dimensional characterization of blood flow in the right ventricle (RV) and right atrium (RA). In this study, we investigate and quantify differences in the characteristics of coherent rotating flow structures (vortices) in the RA and RV between subjects with right ventricular diastolic dysfunction (RVDD) and normal controls. Fifteen RVDD subjects and 10 age-matched controls underwent same day 3D time resolved CMR and echocardiography. Echocardiography was used to determine RVDD stage as well as pulmonary artery systolic pressure (PASP). CMR data was used for RA and RV vortex quantification and visualization during early ventricular diastole and the results are compared between healthy subjects and those with RVDD. The resulting trends are discussed and hypotheses are presented regarding differences in vortex characteristics between healthy and RVDD subjects cohorts.

James Browning
University of Colorado Boulder

Date submitted: 03 Aug 2015

Electronic form version 1.4