

Abstract Submitted
for the DFD16 Meeting of
The American Physical Society

Rotating thermal convection at very large Rayleigh numbers¹

STEPHAN WEISS, Max Planck Institute f. Dynamics and Self-Organisation, DENNIS VAN GILS, University of Twente, GUENTER AHLERS, University of California, Santa Barbara, EBERHARD BODENSCHATZ, Max Planck Institute f. Dynamics and Self-Organisation — The large scale thermal convection systems in geo- and astrophysics are usually influenced by Coriolis forces caused by the rotation of their celestial bodies. To better understand the influence of rotation on the convective flow field and the heat transport at these conditions, we study Rayleigh-Bénard convection, using pressurized sulfur hexafluoride (SF_6) at up to 19 bars in a cylinder of diameter $D=1.12$ m and a height of $L=2.24$ m. The gas is heated from below and cooled from above and the convection cell sits on a rotating table inside a large pressure vessel (the “Uboot of Göttingen”). With this setup Rayleigh numbers of up to $Ra = 10^{15}$ can be reached, while Ekman numbers as low as $Ek = 10^{-8}$ are possible. The Prandtl number in these experiment is kept constant at $Pr \simeq 0.8$. We report on heat flux measurements (expressed by the Nusselt number Nu) as well as measurements from more than 150 temperature probes inside the flow.

¹We thank the Deutsche Forschungsgemeinschaft (DFG) for financial support through SFB963: Astrophysical Flow Instabilities and Turbulence. The work of GA was supported in part by the US National Science Foundation through Grant DMR11-58514.

Stephan Weiss
Max Planck Institute f. Dynamics and Self-Organisation

Date submitted: 23 Jul 2016

Electronic form version 1.4