

Abstract Submitted
for the DFD20 Meeting of
The American Physical Society

Statistical transition to turbulence in plane channel flow

SÉBASTIEN GOMÉ, LAURETTE TUCKERMAN, Laboratoire de Physique et Mécanique des Milieux Hétérogènes, CNRS, ESPCI Paris, PSL, Sorbonne Université, Univ. de Paris, DWIGHT BARKLEY, Mathematics Institute, University of Warwick, Coventry CV4 7AL, United Kingdom — Transition to turbulence in shear flows is characterized by intermittent laminar-turbulent patterns, which statistically proliferate or decay depending on the Reynolds number. The evolution of turbulent bands in plane channel flow is studied via direct numerical simulations in a narrow computational domain tilted with respect to the streamwise direction. Bands interact via their intervening quasi-laminar gap, impacting their propagation velocities. Turbulence decay and spreading processes are in most cases exponentially distributed, which is the signature of a memoryless process. Statistically estimated time scales for band decay or splitting depend super-exponentially on the Reynolds number and lead to the estimation of Reynolds number $Re_{\text{cross}} \simeq 965$ above which splitting is more likely than decay. The associated time scales are over 10^6 advection times. With such a low associated probability, the use of statistical mechanics methods such as a rare event algorithm is necessary to evaluate the mean decay or splitting time. Decay or splitting events are analyzed through the dynamics of large-scale Fourier components of the velocity, which statistically approaches a most-probable pathway.

Sébastien Gomé
Sorbonne Université

Date submitted: 31 Jul 2020

Electronic form version 1.4