

Abstract Submitted  
for the DFD20 Meeting of  
The American Physical Society

**Turbulence explains the accelerations of an eagle in natural flight**

KASEY LAURENT, Cornell University, BOB FOGG, Cellular Tracking Technologies, TOBIAS GINSBURG, Cornell University, CASEY HALVERSON, MIKE LANZONE, Cellular Tracking Technologies, TRICIA MILLER, Conservation Science Global, DAVID WINKLER, Conservation Science Global, SABER Consulting, GREGORY BEWLEY, Cornell University — Soaring birds travel hundreds of miles without flapping their wings, and previous work has shown birds' ability to utilize large-scale atmospheric flow structures such as thermal and orographic updrafts to enhance flight. However, it is unclear what role turbulence plays in avian flight behavior. We analyzed acceleration and GPS data from a golden eagle (*Aquila chrysaetos*) and found that the bird's accelerations can be entirely explained by linear interactions with turbulence within an interval of timescales between about 0.5 and 10 seconds. We isolated soaring flight from other behaviors by finding patterns in the accelerations and by adapting known methods for behavior classification. From the position data, we approximated the wind speeds experienced by the golden eagle, which were between 2 and 13 m/s. We found that the probability distribution function and the spectrum of the eagle's accelerations are non-Gaussian and resemble those for lightly inertial particles in turbulence. Additionally, we find that higher wind speeds are associated with larger accelerations, corresponding to a lifting of the spectra consistent with the increasing strength of turbulence in higher winds.

Kasey Laurent  
Cornell University

Date submitted: 03 Aug 2020

Electronic form version 1.4