## Abstract Submitted for the DNP06 Meeting of The American Physical Society

Coincidence Geometry Measurements of the  ${}^{1}S_{0}$  Scattering Lengths in Neutron-Deuteron Breakup at 19  $MeV^1$  A.S. CROWELL, J. DENG, J.H. ESTERLINE, C.R. HOWELL, M.R. KISER, R.A. MACRI, S. TAJIMA, W. TORNOW, Duke University and TUNL, B.J. CROWE III, North Carolina Central University, R.S. PEDRONI, North Carolina A&T State University, W. VON WITSCH, University of Bonn, H. WITAŁA, Jagellonian University — Measurements of the  ${}^{1}S_{0}$  neutron-neutron (nn) and neutron-proton (np) scattering lengths,  $a_{nn}$  and  $a_{np}$  respectively, using neutron-induced deuteron breakup were made at Triangle Universities Nuclear Laboratory (TUNL) at an incident neutron energy of 19.0 MeV. Six liquid scintillator detectors were configured in a coincidence geometry to measure the momenta of the two neutrons in two nn and np finalstate-interaction (FSI) pairs while the energy of the proton was determined using a  $C_6D_{12}$  active target. The scattering lengths were extracted from the experimental cross sections by comparison to rigorous three-nucleon calculations using the CD Bonn nucleon-nucleon potential for various values of  $a_{nn}$  and  $a_{np}$ . In this talk results from the two nn and np FSI measurements and the analysis to obtain  $a_{nn}$  will be presented.

<sup>1</sup>This work was supported in part by USDOE Grant No. DE-FG02-97-ER41033.

Alexander Crowell Duke University and TUNL

Date submitted: 30 Jun 2006

Electronic form version 1.4