Abstract Submitted for the DNP06 Meeting of The American Physical Society

In-beam gamma-ray spectroscopy of $172Os^1$ K. ALEKSANDROVA, P. MANCHEV, M.S. FETEA, Department of Physics, University of Richmond, A. HEINZ, WNSL, Yale University, G. GURDAL, WNSL, Yale University and Clark University, R. CASPERSON, R.F. CASTEN, WNSL, Yale University, M. CHAMBERLAIN, University of Surrey, UK and WNSL, Yale University, E.A. MC-CUTCHAN, J. QIAN, WNSL, Yale University, N.J. THOMPSON, University of Surrey, UK and WNSL, Yale University, V. WERNER, R. WINKLER, WNSL, Yale University — A 30Si beam was used for a test experiment at the WNSL facility. The production of the compound nucleus 174Os was achieved through complete fusion in a 100 µg/cm2 144Sm target at beam energy of 134MeV. Gamma-rays from residual nuclei were detected with the YRAST ball array consisting of 7 Comptonsuppressed clover detectors. A number of Os isotopes, in particular 172Os were also detected. While the main goal of the experiment was the commissioning of a gas-filled detector, we present here data from in-beam gamma-ray spectroscopy.

¹This work was supported by NSF grants PHY 0204811 and 0555665, Jeffress Fund grant J-809, U.S. DOE Grant Nos. DE-FG02-91ER-40609, DE-FG02-88ER-40417, DE-FG02-05ER40417.

Kalina Aleksandrova Department of Physics, University of Richmond

Date submitted: 01 Aug 2006

Electronic form version 1.4