

Abstract Submitted  
for the DNP08 Meeting of  
The American Physical Society

**Track Reconstruction for the NIFFTE TPC** SARVAGYA SHARMA, Abilene Christian University, NEUTRON INDUCED FISSION FRAGMENT TRACKING EXPERIMENT COLLABORATION — The Global Nuclear Energy Partnership (GNEP) has funded the construction of a Time Projection Chamber (TPC) to be used for precision fission cross-section measurements through the Nuclear Energy Research Initiative (NERI). This poster shall illustrate the status of algorithms intended for intelligent track finding and track fitting using raw data obtained TPC simulations will be presented. The track fitting effort in this experiment has borrowed a number of ideas from high-energy physics along with other pattern recognition techniques not previously affiliated with experimental physics. Two track-finding techniques have been investigated. The Hough Transform is a brute force attempt at finding tracks. The second paradigm for track reconstruction, Binary Space Partitioning (BSP), was found to be less computationally expensive. BSP however had to be used in conjunction with a multilayer perceptron to rejoin tracks that were overly partitioned. To determine track fit parameters, an iterative Kalman Filter has been implemented that allows multiple scattering and energy losses to be taken into account to obtain unbiased errors.

Sarvagya Sharma  
Abilene Christian University

Date submitted: 14 Aug 2008

Electronic form version 1.4