

Abstract Submitted  
for the DNP10 Meeting of  
The American Physical Society

**A measurement of the differential cross section for the reaction  $\gamma^*n \rightarrow \pi^-p$  from deuterium** JIXIE ZHANG, Jefferson Lab, Newport News, VA, 23606, GAIL DODGE, Old Dominion University, Norfolk, VA, 23529, THE CLAS COLLABORATION — Differential cross sections for exclusive  $\pi^-$  electroproduction from the neutron in the reaction  $e+d \rightarrow e'+\pi^-+p+p_r$  have been measured over a broad kinematics range. The experiment was performed using a newly built radial time projection chamber (RTPC) and the CEBAF Large Acceptance Spectrometer (CLAS) at Jefferson Lab. The RTPC detector was specially built to detect low energy recoil protons and had a valid momentum acceptance from 67 to 250 MeV/c. Electron scattering data were taken with beam energies of 2.1, 4.2 and 5.3 GeV using a 7 atm gaseous deuterium target. The differential cross sections for  $D(e, e'\pi^-p)p$  have been extracted, with the proton detected either by the CLAS or by the RTPC. The structure functions  $\sigma_T + \varepsilon\sigma_L$ ,  $\sigma_{LT}$  and  $\sigma_{TT}$  are also extracted and compared to MAID and SAID predictions.

Jixie Zhang  
Jefferson Lab, Newport News, VA, 23606

Date submitted: 01 Jul 2010

Electronic form version 1.4