

Abstract Submitted
for the DNP12 Meeting of
The American Physical Society

Spectroscopy of neutron-unbound ^{15}Be JESSE SNYDER, MICHAEL THOENNESSEN, THOMAS BAUMANN, ARTEMIS SPYROU, MICHAEL STRONGMAN, GREG CHRISTIAN, SHEA MOSBY, MICHELLE MOSBY, JENNA SMITH, ANNA SIMON, NSCL/MSU, BRYAN LUTHER, Concordia College, SHARON STEPHENSON, ALEX PETERS, Gettysburg College, PAUL DEYOUNG, ERIC LUNDERBERG, Hope College, JOSEPH FINCK, Central Michigan University — A (d,p) reaction was used to populate neutron unbound states in ^{15}Be from a secondary beam of ^{14}Be at 55 MeV/u impinging upon a deuterated polyethylene target. The unbound states in ^{15}Be decayed through the emission of a neutron. The neutron was detected using MoNA, an array of 144 plastic scintillator bars. The charged ^{14}Be fragments were deflected by the Sweeper dipole magnet into a system of charged particle detectors which allow the reconstruction of their kinematic properties. The decay energy was calculated through invariant mass analysis, using the energy and momentum information of the neutron and fragment at the target. Results will be presented.

Jesse Snyder
NSCL/MSU

Date submitted: 14 Aug 2012

Electronic form version 1.4