

Abstract Submitted  
for the DNP12 Meeting of  
The American Physical Society

**Fixed Target Results from STAR using Gold on Beampipe (Al) Events** CHRISTOPHER FLORES, DANIEL CEBRA, SAMANTHA BROVKO, University of California at Davis, BROOKE HAAG, Hartnell College, JIM DRAPER, University of California at Davis, STAR COLLABORATION — The RHIC Beam Energy Scan was proposed to search for the possible critical point and to study the nature of the phase transition between hadronic and partonic matter. However, several dynamical model simulations (UrQMD, PHSD, QGSM, GiBUU, 3-fluid) suggest that the partonic phase is entered for center-of-mass collision energies as low as 4-5 GeV [1]. Collisions between beam halo nuclei and the aluminum beam pipe allow STAR to study fixed-target Au+Al collisions. The injection and sub-injection energy gold beams (kinetic energies of 8.8, 4.8 and 2.9 AGeV) produce Au+Al collisions at center-of-mass energies of 4.5, 3.5, and 3.0 GeV. Particle ratios will be presented and compared to earlier published results from the AGS. Fixed target acceptances and efficiencies for tracking in the TPC and particle identification in the Time of Flight system will be shown.

[1] I. C. Arsene et al., Phys. Rev. C 75, 034902 (2007)

Christopher Flores  
University of California at Davis

Date submitted: 06 Jul 2012

Electronic form version 1.4