

Abstract Submitted  
for the DNP12 Meeting of  
The American Physical Society

**Study of light tin isotopes via single-nucleon knockout reactions<sup>1</sup>**

A. AYRES, K.L. JONES, University of Tennessee, T. BAUGHER, D. BAZIN, J. BERRYMAN, NSCL, A. BEY, C. BINGHAM, L. CARTEGNI, G. CERIZZA, University of Tennessee, A. CHAE, Sungkyunkwan University/ORNL, A. GADE, NSCL, R. GRZYWACZ, University of Tennessee, M.E. HOWARD, Rutgers University, S. MCDANIEL, NSCL, D. MILLER, University of Tennessee/TRIUMF, S. PADGETT, University of Tennessee/LLNL, S. PAIN, ORNL, A. RATKIEWICZ, NSCL/Rutgers University, A. SHORE, R. STROBERG, D. WEISSHAAR, K. WIMMER, R. WINKLER, NSCL — Spectroscopic studies have been performed close to  $^{100}\text{Sn}$ , utilizing the S800 and CAESAR at the NSCL. These studies make use of a single neutron knockout reaction on  $^{108}\text{Sn}$  and  $^{106}\text{Sn}$  beams. The momentum distributions of the resulting residues indicate the  $\ell$ -value of the removed neutron and the spectroscopic factors for the even-mass nuclei. Additionally,  $\gamma$ -rays were measured in coincidence with the momentum distributions allowing the separation of the knockout channel where the residue is left in an excited state from the channel to the ground state. The odd-mass residue can then be characterized in terms of a hole in the d- or g- orbital with reference to the even-mass nucleus. The current status of the analysis will be presented.

<sup>1</sup>Research supported in part by the NSF, U.S. DOE Office of Nuclear Physics, and the NNSA.

A. Ayres  
University of Tennessee

Date submitted: 10 Jul 2012

Electronic form version 1.4