

Abstract Submitted
for the DNP15 Meeting of
The American Physical Society

Calibrations for studies of neutron-rich precursor fragments¹

MARIA MAZZA, Gettysburg College, RACHEL PARKHURST, Westmont College, SAMUEL WILENSKY, Gettysburg College, MICHELLE MOSBY, LANL, SHARON STEPHENSON, Gettysburg College, WARREN ROGERS, Westmont College, MONA COLLABORATION COLLABORATION — Heavy ion collisions at relativistic energies produce the radioactive beams used at nuclear structure facilities worldwide. However, there are still unanswered questions about the reaction mechanism of projectile fragmentation and the specific roles that ablation, evaporation, and abrasion play. Using the projectile fragmentation of a ^{32}Mg beam at 86 MeV/u on a natural Beryllium target at the National Superconducting Cyclotron Laboratory (NSCL), our experimental goal is to better understand the excitation energy and the momentum distribution of the precursors of the observed final fragments (neon, sodium, and fluorine). A suite of charged particle detectors in conjunction with the Modular Neutron Array (MoNA) allows us to analyze both the charged final fragments as well as the coincident neutrons. Detector calibration results and preliminary results will be presented.

¹This material is based upon work supported by the National Science Foundation under grants PHY-1205537.

Sharon Stephenson
Gettysburg College

Date submitted: 31 Jul 2015

Electronic form version 1.4