

Abstract Submitted
for the DNP16 Meeting of
The American Physical Society

A high-precision measurement of the $\pi \rightarrow e\nu$ branching ratio

TRISTAN SULLIVAN, D.A. BRYMAN, S. CUEN-ROCHIN, University of British Columbia, A. AGUILAR-AREVALO, Universidad Nacional Autónoma de Mexico, M. AOKI, Osaka University, M. BLECHER, Virginia Tech, D.I. BRITTON, D. PROTOPODESCU, University of Glasgow, D. VOM BRUCH, Johannes Gutenberg-University Mainz, S. CHEN, Tsinghua University, J. COMFORT, Arizona State University, L. DORIA, P. GUMPLINGER, L. KURCHANINOV, R.E. MISCHKE, T. NUMAO, A. SHER, D. VAVILOV, TRIUMF, A. HUSSEIN, University of Northern British Columbia, Y. IGARASHI, . KEK, S. ITO, Okayama University, S.H. KETTELL, L.S. LITTENBERG, Brookhaven National Laboratory, C. MALBRUNOT, CERN — The pion decay branching ratio, $R_\pi = \frac{\Gamma(\pi \rightarrow e\nu + \pi \rightarrow e\nu\gamma)}{\Gamma(\pi \rightarrow \mu\nu + \pi \rightarrow \mu\nu\gamma)}$, is an important observable in the Standard Model of particle physics. The value of the $\pi \rightarrow e\nu$ branching ratio has been calculated within the Standard Model to be $(1.2352 \pm 0.0002) \times 10^{-4}$. The PIENU experiment at TRIUMF aims to measure this quantity to a precision of $< 0.1\%$. This provides a sensitive test of lepton universality, and constraints on many new physics scenarios with mass reach up to 1000 TeV. The measurements are also sensitive to the presence of heavy neutrinos. The current state of the analysis of the PIENU data will be presented.

TRISTAN SULLIVAN
University of British Columbia

Date submitted: 06 Oct 2016

Electronic form version 1.4