

Abstract Submitted
for the DNP17 Meeting of
The American Physical Society

Search for the permanent electric dipole moment of ^{129}Xe

NATASHA SACHDEVA, TIMOTHY CHUPP, FEI GONG, University of Michigan, USA, EARL BABCOCK, ZAHIR SALHI, Jülich Centre for Neutron Science, Germany, MARTIN BURGHOFF, ISAAC FAN, WOLFGANG KILLIAN, SILVIA KNAPPE-GRÜNEBERG, ALLARD SCHABEL, FRANK SEIFERT, LUTZ TRAHMS, JENS VOIGT, Physikalisch-Technische Bundesanstalt, Germany, SKYLER DEGENKOLB, Institut Laue-Langevin, France, PETER FIERLINGER, EVA KRÄGELOH, TOBIAS LINS, MICHAEL MARINO, JONAS MEINEL, BEN-JAMIN NIESSEN, STEFAN STUIBER, WILLIAM TERRANO, Excellence Cluster Universe and Technische Universität München, Germany, FLORIAN KUCHLER, TRIUMF, Canada, JAIDEEP SINGH, Michigan State University and NSCL, USA — CP-violation in Beyond-the-Standard-Model physics, necessary to explain the baryon asymmetry, gives rise to permanent electric dipole moments (EDMs). EDM measurements of the neutron, electron, paramagnetic and diamagnetic atoms constrain CP-violating parameters. The current limit for the ^{129}Xe EDM is $6 \times 10^{-27} e \cdot \text{cm}$ (95% CL). The HeXeEDM experiment at FRM-II (Munich Research Reactor) and BMSR-2 (Berlin Magnetically Shielded Room) uses a stable magnetic field in a magnetically shielded room and ^3He comagnetometer with potential to improve the limit by two orders of magnitude. Polarized ^3He and ^{129}Xe free precession is detected with SQUID magnetometers in the presence of applied electric and magnetic fields. Conclusions from recent measurements will be presented.

Natasha Sachdeva
Univ of Michigan - Ann Arbor

Date submitted: 30 Jun 2017

Electronic form version 1.4