

Abstract Submitted
for the DNP19 Meeting of
The American Physical Society

Constraining Sodium Production in Globular Clusters Using the $^{23}\text{Na}(^3\text{He}, d)^{24}\text{Mg}$ Reaction¹ CALEB MARSHALL, North Carolina State University, TUNL, KIANA SETOODEHNIA, European X-ray Free Electron Laser GmbH, FEDERICO PORTILLO, RICHARD LONGLAND, North Carolina State University, TUNL — Globular clusters consist of hundreds of thousands of stars gravitationally bound in a relatively small radius (~ 10 pc). Over the last few decades, intense observational study has revealed that globular clusters are comprised of multiple stellar populations each with distinct chemical signatures.² The star-to-star Na-O anticorrelation is the most pervasive of these so called abundance anomalies, and is theorized to be the result of stellar material undergoing hydrogen burning at 50–100 MK. Unfortunately, many thermonuclear reaction rates suffer from large uncertainties at these temperatures, thereby limiting our understanding of nucleosynthesis in globular clusters. Among these rates one of the most critical is the sodium destroying reaction $^{23}\text{Na}(p, \gamma)^{24}\text{Mg}$. Using the Enge Split-pole Spectrograph at Triangle Universities Nuclear Laboratory (TUNL), we have measured the transfer reaction $^{23}\text{Na}(^3\text{He}, d)^{24}\text{Mg}$. Excitation energies, spin-parities, and spectroscopic factors were extracted for states of interest improving our estimates of the $^{23}\text{Na}(p, \gamma)$ reaction rate, and thus constraining sodium destruction in stellar material.

¹This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Award Numbers DE-SC0017799 and under Contract No. DE-FG02-97ER41041.

²R. Gatton, E. Carretta, A. Bragaglia, **AAPR** 20, 2012

Caleb Marshall
North Carolina State University

Date submitted: 28 Jun 2019

Electronic form version 1.4