

Abstract Submitted
for the DNP19 Meeting of
The American Physical Society

Charged Particle Distributions in Central Au+Au Collisions at $\sqrt{s_{NN}} = 3.0$ GeV at STAR¹ BENJAMIN KIMELMAN, University of California, Davis, STAR COLLABORATION — The RHIC Beam Energy Scan phase I (BES-I) program provided a detailed study of nuclear matter over a wide range of energies. Below $\sqrt{s_{NN}} < 19.6$ GeV, interesting results were shown in hadron azimuthal anisotropies, particle ratios, and net-proton higher moments. These results motivate the Beam Energy Scan phase II (BES-II). Compared to BES-I, BES-II will have improvements including increased statistics by a factor of 10 to 20 for each energy, improved acceptance from upgrades to the STAR experiment, and an extension of the energy reach from $\sqrt{s_{NN}} = 7.7$ GeV to $\sqrt{s_{NN}} = 3.0$ GeV with the STAR fixed-target program. This talk will present results from the lowest fixed target energy to be studied in BES-II including transverse mass spectra, rapidity density distributions, particle ratios, and centrality dependence for charged hadrons. These results are analyzed with a chemical equilibrium model to determine the chemical temperature and potential at freeze-out. The pion ratio and Coulomb potential will also be presented. At low energy, produced particles are sensitive to a Coulomb potential from a net positive source at low momenta which modifies the transverse mass spectra. These new data are compared to previously published results from experiments at the AGS.

¹NSF Grant No. 1812398

Benjamin Kimelman
University of California, Davis

Date submitted: 30 Jun 2019

Electronic form version 1.4