

Abstract Submitted
for the DNP20 Meeting of
The American Physical Society

ϕ -meson production in Au + Au collisions at $\sqrt{s_{\text{NN}}} = 3$ GeV from STAR GUANNAN XIE, Lawrence Berkeley National Laboratory, STAR COLLABORATION — ϕ -meson is the lightest bound state of strange quarks($s\bar{s}$). It has relatively small hadronic interaction cross sections, therefore ϕ -meson is considered to be a sensitive probe of the earlier dynamics in the heavy-ion collision. Recent measurements by HADES and FOPI on subthreshold ϕ -meson production show a larger ϕ/K^- ratio compared to the results at higher energies[1,2], and this larger ϕ/K^- ratio cannot be described by thermal model calculations with Grand Canonical Ensemble for strangeness. In this presentation, we will report on our first measurements of ϕ -meson production in Au+Au collisions at $\sqrt{s_{\text{NN}}} = 3$ GeV with energy just above the NN threshold. The data were taken in 2018 by the STAR experiment with Fixed Target configuration. ϕ -mesons are measured through their hadronic decay channel, $\phi \rightarrow K^- + K^+$. After being corrected for the detector acceptance and tracking efficiencies, invariant yields of ϕ -mesons as well as ϕ/K^- ratio are presented in several centrality intervals and the results will be compared to model calculations.

HADES Collaboration, Phys. Let. **B 778**, (2018) 403-407.

FOPI Collaboration, Eur. Phys. J. **A 52**, (2016) 177.

Guannan Xie
Lawrence Berkeley National Laboratory

Date submitted: 24 Jun 2020

Electronic form version 1.4