Abstract Submitted for the DPP05 Meeting of The American Physical Society

**Comparison of DIII-D Phase Contrast Imaging Measurements in** the Edge and Core Regions<sup>1</sup> C.J. ROST, M. PORKOLAB, J.R. DORRIS, Massachusetts Institute of Technology, K.H. BURRELL, General Atomics — During the last DIII-D run period, the phase contrast imaging (PCI) turbulence diagnostic acquired data in an upgraded configuration. The improvements include 10 MHz digitizers with a data record covering the entire discharge and a wavenumber range increased to 30/cm. The beampath was previously tangent to the LCFS but now passes through the LCFS and reaches r/a=0.8. The PCI was previously sensitive only to radial modes, but it is now sensitive to modes with finite poloidal wavenumber. Measurements of turbulence near the ITG range, particularly the S(k,f) spectra, now show a Doppler shift that was never observed previously. Analysis of this new data includes the variation of the magnetic field along the beam path. Comparisons with previous PCI measurements in the old beam geometry give us a more complete picture of the edge turbulence. The increased wavenumber range also allows us to examine how experimental frequency spectra of plasma turbulence depend on the wavenumber range of the diagnostic.

<sup>1</sup>Work supported by US DOE under DE-FG02-94ER54235 and DE-FC02-04ER54698.

T.S. Taylor General Atomics

Date submitted: 21 Jul 2005

Electronic form version 1.4