

Abstract Submitted
for the DPP05 Meeting of
The American Physical Society

Particle simulation of a virtual cathode in bipolar flow¹ JOHN VERBONCOEUR, University of California-Berkeley, JEAN-LUC VAY, Lawrence Livermore National Laboratory, PETER STOLTZ, TechX Corp. — Virtual cathode formation is studied in a bipolar flow relevant to the High Current Experiment parameter regime. A 174 mA potassium ion beam with beam energy of 972 keV, confined in a conducting drift tube, is incident on a stainless steel plate. The ion impacts generate secondary electrons with an energy and angular dependent yield large compared to 1, resulting in a current sufficient for formation of a virtual cathode. Using enhanced ion-induced secondary models in the PIC codes XOOPIE and WARP, the formation of the virtual cathode and its effect on the bipolar flow is investigated, including time-dependent effects. Numerical issues including resolution of the virtual cathode scale length $l_{vc} \sim \lambda_{De}$ and time scale $t \sim 1/\omega_{pe}$ are characterized. Results indicate an improvement in temporal and spatial fidelity leads to a lower and more stable virtual cathode potential as well as significant impact on neutrality in the bipolar flow.

¹This work was supported in part under the auspices of the U.S. Department of Energy by University of California, LLNL, and LBL under contracts W-7405-Eng-48 and DE-AC03-76F00098, and in part by the Air Force Research Laboratory-Kirtland.

John Verboncoeur
University of California-Berkeley

Date submitted: 27 Jul 2005

Electronic form version 1.4