

Abstract Submitted
for the DPP06 Meeting of
The American Physical Society

Heavy ion beam driven warm dense matter experiments¹ F.M. BIENIOSEK, M.A. LEITNER, B.G. LOGAN, R.M. MORE, P.K. ROY, LBNL, J.J. BARNARD, LLNL, L.R. GRISHAM, PPPL — We describe near term heavy-ion beam-driven warm dense matter (WDM) experiments. Initial experiments are at low beam velocity, below the Bragg peak, increasing toward the Bragg peak in subsequent versions of the accelerator. The WDM conditions are envisioned to be achieved by combined longitudinal and transverse neutralized drift compression to provide a hot spot on the target with a beam spot size of about 1 mm, and pulse length about 1-2 ns. The range of the beams in solid matter targets is about 1 micron, which can be lengthened by using porous targets at reduced density. Initial candidate experiments include an experiment to study transient darkening in the WDM regime; and a thin target dE/dx experiment to study beam energy and charge state distribution in a heated target. Further experiments will explore target temperature and other properties such as electrical conductivity to investigate phase transitions and the critical point.

¹This work performed under the auspices of the U.S Department of Energy by Lawrence Livermore and Lawrence Berkeley National Laboratories and Princeton Plasma Physics Laboratory under contracts No. W-7405-Eng-48, DE-AC02-05CH11231, and DE-AC02-76CH03073.

F. M. Bieniosek
LBNL

Date submitted: 07 Jul 2006

Electronic form version 1.4