

Abstract Submitted
for the DPP06 Meeting of
The American Physical Society

Fast Ion Generation and Confinement during Reconnection Events in the MST Reversed Field Pinch¹ RICHARD MAGEE, BRETT CHAPMAN, GENNADY FIKSEL, DARREN CRAIG, DAVID ENNIS, SANJAY GANGADHARA, University of Wisconsin - Madison, MADISON SYMMETRIC TORUS TEAM — Measurements of neutron flux from deuterium plasmas in the MST reversed field pinch used in conjunction with ion temperature measurements indicate the presence of a fast ion population generated at magnetic reconnection events. During a typical event, T_i on-axis approximately doubles, from ~ 0.5 keV to ~ 1 keV, in less than $200 \mu\text{s}$. These events often produce neutron fluxes up to ten times higher than from thermal fusion alone. This flux is consistent with, for example, a small (1%), non-thermal population at ~ 15 keV. After an event, the neutron flux decays with a time constant of 1 - 3 ms. However, if an event is followed by a period of reduced magnetic fluctuations (achieved either actively, by inductively driving a parallel current in the edge, or spontaneously), the neutron flux decays at a much slower rate, ~ 20 ms, which indicates improved fast ion confinement.

¹Work supported by US DoE and NSF

Richard Magee
Department of Physics, University of Wisconsin - Madison

Date submitted: 23 Aug 2006

Electronic form version 1.4