

Abstract Submitted  
for the DPP06 Meeting of  
The American Physical Society

**Experiments at High  $I_p/I_{tf}$  in the Pegasus Toroidal Experiment<sup>1</sup>**

E.A. UNTERBERG, D.J. BATTAGLIA, M.W. BONGARD, N.W. EIDIETIS, R.J. FONCK, M.J. FROST, G.D. GARSTKA, B.J. SQUIRES, M.B. MCGARRY, University of Wisconsin-Madison — The operating space defined by the external kink mode boundary in a near-unity aspect ratio ST allows access to very high toroidal beta and  $I_N$  through operations at high toroidal field utilization factors,  $I_p/I_{tf} > 1$ . In Pegasus, however, purely inductive-driven plasmas exhibit an operational limit of  $I_p/I_{tf} \sim 1$  due to the onset of large-scale tearing modes in a wide volume of low shear near the plasma core region. A new operating regime has been accessed in recent experiments with the addition of electrostatic helicity and edge-current sources. Employing these sources in the lower divertor region as a pre-ionization source, inductive startup at very low field is facilitated and allows  $I_p/I_{tf} \approx 1.5$  at low  $I_p$  ( $\sim 50$ kA). Using these sources for non-inductive startup via helicity injection provides access to  $I_p/I_{tf} \approx 2.3$ , again at low  $I_p$ . In both cases, no large-scale tearing modes are evident. These observations, coupled with magnetic reconstructions, indicating hollow  $j(R)$  with possible reverse shear, suggest the sources provide significant modifications to the  $j(R)$  profile to allow stable discharge evolution at minimal toroidal field. Experimental studies are focusing on extending these recent discharges to higher  $I_p$ .

<sup>1</sup>Work supported by the U.S. D.O.E. Grant DE-FG02-96ER54375

Gregory Garstka  
University of Wisconsin-Madison

Date submitted: 21 Jul 2006

Electronic form version 1.4