

Abstract Submitted  
for the DPP08 Meeting of  
The American Physical Society

**The effect of plasma triangularity on turbulent transport: modelling TCV experiments by linear and non linear gyrokinetic simulations**

ALESSANDRO MARINONI, STEPHAN BRUNNER, YANN CAMENEN, STEFANO CODA, JONATHA GRAVES, XAVIER LAPILLONNE, ANTOINE POCH-ELON, OLIVIER SAUTER, LAURENT VILLARD, Ecole Polytechnique Federale de Lausanne (EPFL), Centre de Recherches en Physique des Plasmas CH-1015 Lausanne, Association Euratom-Conf. Suisse, ECOLE POLYTECHNIQUE FEDERALE DE LAUSANNE (EPFL), CENTRE DE RECHERCHES EN PHYSIQUE DES PLASMAS TEAM — Experimental evidence in the TCV tokamak revealed that the core electron heat transport is significantly reduced by a negative triangularity ( $\delta$ ) configuration. This variation in heat transport cannot be explained by the flux surfaces compression; rather, it must be due to a (partial) stabilization of the microinstabilities at play in a positive  $d$  configuration. The present work is a theoretical investigation of the effect exerted by triangularity on plasma turbulence of actual TCV shots, using the gyrokinetic code GS2. A complex interplay of deeply and barely trapped particles, which react in opposite ways to triangularity, results in a stabilizing effect of negative  $\delta$  on Trapped Electron Modes both in the linear and non-linear phases. The non-linear simulations are in qualitative agreement with the experimental results.

Stefano Coda  
Ecole Polytechnique Federale de Lausanne (EPFL),  
Centre de Recherches en Physique des Plasmas  
CH-1015 Lausanne, Association Euratom-Conf. Suisse

Date submitted: 01 Aug 2008

Electronic form version 1.4