

Abstract Submitted  
for the DPP08 Meeting of  
The American Physical Society

**3D Modeling of the Interaction of the Solar Wind with the Interstellar Medium** GARY ZANK, University of Alabama at Huntsville — The presence of the interplanetary magnetic field and current sheet and the interstellar magnetic field render the interaction of the solar wind with the local interstellar medium intrinsically 3D. The inclusion of magnetic fields into the problem is therefore challenging from a simulation perspective. Furthermore, the solar wind is itself intrinsically time-dependent on a range of time scales and one has to ideally model the interaction as time variable. Finally, to further complicate matters, interstellar neutral gas, primarily H, is coupled to both LISM and interplanetary plasma through charge exchange and it has been shown in the context of simpler 2D models that neutral H can modify the global structure of the heliosphere profoundly. We discuss here the extension of earlier 2D models of the solar wind-LISM interaction to include interplanetary and interstellar magnetic fields, i.e., modeling the 3D heliosphere. We will demonstrate the profoundly important role that neutral H continues to play in determining the structure of the global heliosphere, and the corresponding back-reaction on the neutral H distribution. This will be addressed in the context of both multi-fluid and kinetic models that have been developed by our group. We will discuss the origin of energetic neutral atoms using an approach based on a kappa distribution description of interplanetary plasma.

Gary Zank  
University of Alabama at Huntsville

Date submitted: 17 Jul 2008

Electronic form version 1.4