

Abstract Submitted
for the DPP08 Meeting of
The American Physical Society

Analytic and computational investigation of the effect of finite parallel heat transport on the formation of magnetic islands in 3-D plasma equilibria¹ M. SCHLUTT, C.C. HEGNA, Univ. of Wisconsin, E. HELD, Utah State Univ. — A resistive MHD model is used to investigate pressure-induced magnetic islands in 3-D equilibria. We revisit previous island calculations while allowing for finite parallel heat transport, to derive an equation for equilibrium island width. We find that finite parallel heat transport can alter the impact of resistive interchange and bootstrap current contributions to magnetic island formation. However, Pfirsch-Schlüter currents driven by resonant components in $\frac{1}{B^2}$ are largely unaffected by transport processes. To make further progress, this problem is modeled using NIMROD. A vacuum equilibrium helical magnetic field is loaded into the geometry of a straight stellarator. The symmetry of the vacuum field with a dominant magnetic harmonic can be spoiled by adding small magnetic perturbations. These perturbations alter the magnetic spectrum, and potentially produce magnetic islands and regions of stochasticity. Numerical simulations will be performed that include the effect of a heating source and self-consistent anisotropic transport processes in a variety of magnetic configurations. The effect of finite parallel heat transport on the island formation and saturation processes will be examined.

¹Supported by U.S. DOE Grant No. DE-FG02-99ER54546.

M. Schlutt
Univ. of Wisconsin

Date submitted: 17 Jul 2008

Electronic form version 1.4