

Abstract Submitted
for the DPP08 Meeting of
The American Physical Society

Multi-cm Long Magnetically Controlled Optical Plasma Channel

BRADLEY POLLOCK, University of California, San Diego & Lawrence Livermore National Laboratory, DUSTIN FROULA, Lawrence Livermore National Laboratory, GEORGE TYNAN, University of California, San Diego, LAURENT DIVOL, PAUL DAVIS, JOHN PALASTRO, DWIGHT PRICE, SIEGFRIED GLENZER, Lawrence Livermore National Laboratory — Experimental results from the Jupiter Laser Facility at the Lawrence Livermore National Laboratory which show multi-cm long plasma channels with minimum plasma densities below $5 \times 10^{17} \text{ cm}^{-3}$ are presented. These results are obtained using an external magnetic field (< 5 Tesla) to limit the radial heat flux from a pre-forming laser beam. The resulting increased plasma pressure gradient produces a parabolic density gradient which is shown to be tunable by changing the external magnetic field strength. These results are compared with 3-D resistive MHD modeling. For these channel conditions, quasi-static kinetic simulations show that 90% of the energy in a 150 TW short pulse beam is guided over 5 cm and predict electron energy gains of 3 GeV from Laser Wake-field Acceleration. This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344 and was partially funded by the Laboratory Directed Research and Development Program under project tracking code 06-ERD-056.

Bradley Pollock
University of California, San Diego & Lawrence Livermore National Laboratory

Date submitted: 21 Jul 2008

Electronic form version 1.4