

Abstract Submitted
for the DPP09 Meeting of
The American Physical Society

Potential of FAIR at Darmstadt and LHC at CERN for High Energy Density Physics Research: the HEDgeHOB Collaboration

NAEEM A. TAHIR, GSI Darmstadt, ALEXANDER SHUTOV, IGOR LOMONOSOV, IPCP Chernogolovka, A.R. PIRIZ, UCLM Ciudad Real, RUEDIGER SCHMIDT, CERN, DIETER H.H. HOFFMANN, TU Darmstadt, CLAUDE DEUTSCH, LPGP University of Paris — Substantial progress in the development of technology of high quality, well focused, strongly bunched intense particle beams have led to development of a novel, very efficient technique of studying High Energy Density Matter (HEDM) in the laboratory. This method involves generation of large samples of HEDM by isochoric and uniform heating of solid targets by these ion beams. Two huge accelerator projects are on the way in Europe. One is the Facility for Antiprotons and Ion Research (FAIR), at Darmstadt and the other is the Large Hadron Collider (LHC) at CERN. Extensive theoretical work has been carried out over the past years to assess the potential of these accelerators to generate HEDM and several experimental schemes have been proposed [1-4]. A brief overview of this work is presented in this talk. [1] N.A. Tahir et al., PRL 95 (2005) 035001. [2] N.A. Tahir et al., PRL 94 (2005) 135004. [3] N.A. Tahir et al., Nucl. Instr. Meth. A 577 (2007) 238. [4] N.A. tahir et al., PRE 79 (2009) in print.

Naeem A. Tahir
GSI Darmstadt

Date submitted: 27 Apr 2009

Electronic form version 1.4