

Abstract Submitted
for the DPP09 Meeting of
The American Physical Society

Strapping field profile to reproduce transition from slow rise to fast eruptions¹ BAO NGUYEN QUOC HA, PAUL BELLAN, California Institute of Technology — In solar coronal loop experiments at Caltech, the hoop force causes plasma loops to expand unless additional forces are applied. By applying a strapping magnetic field of proper polarity, Hansen and Bellan [1] slowed and even inhibited this expansion. Kliem and Torok [2] predicted that a transition from slow rise to fast eruption occurs if the magnetic field's rate of decay with increasing altitude meets an instability criterion. If the restoring force due to the strapping field decays faster than the hoop force, then the plasma will move from a region where its expansion is decelerated to one where its expansion is accelerated. We have calculated magnetic profiles which attain the instability criterion within the length scale of the Caltech experiment and are constructing an auxiliary coil with independent power supply designed to match the calculated profiles.

[1] J. F. Hansen and P. M. Bellan, *Astrophys. J. Lett.* **563**, L183 (2001)

[2] B. Kleim and T. Torok, *Phys. Rev. Lett.* **96**, 255002 (2006)

¹Work supported by the NSF and AFOSR.

Bao Nguyen Quoc Ha
California Institute of Technology

Date submitted: 08 Jul 2009

Electronic form version 1.4