

Abstract Submitted
for the DPP09 Meeting of
The American Physical Society

Relationship of SOL turbulence to SOL width in Alcator C-Mod¹

STEWART ZWEBEN, PPPL, JAMES TERRY, BRIAN LABOMBARD, MIT, DAREN STOTLER, D.P. LUNDBERG, PPPL, R.H. COHEN, M. UMANSKY, LLNL, D.A. BAVER, D.A. D'IPPOLITO, J.R. MYRA, D.A. RUSSELL, Lodestar — Measurements of SOL turbulence in Alcator C-Mod show a strong variation with poloidal angle [1], therefore it is likely this variation will affect the SOL width. The poloidal variation of the SOL turbulence will be measured as a function of the SOL collisionality, plasma shape, and normalized gyroradius, and the results will be compared with theoretical models for these variations. The radial transport due to the SOL turbulence will be estimated from the radial propagation and correlation length at each poloidal angle, and the results will be compared with the local temperature/density SOL widths measured by Langmuir probes and D_α emission. The expected relationship between these local SOL widths and the total heat flux SOL width at the divertor plate will be discussed.

[1] J.L. Terry, S.J. Zweben et al, J. Nucl. Mat. 390-391, 339 (2009)

¹This work was supported by Work supported by US DOE Contracts #DE-AC02-09CH11466 and DE-FC02-99ER54512.

Stewart Zweben
PPPL

Date submitted: 16 Jul 2009

Electronic form version 1.4