

Abstract Submitted
for the DPP09 Meeting of
The American Physical Society

Transport properties of Hydrogen and CH under ICF conditions

JEAN CLEROUIN, VANINA RECOULES, CHARLES STARRETT, STEPHANE MAZEVET, FLAVIEN LAMBERT, BENOIT CANAUD, ALAIN DECOSTER, CEA/DAM/DIF 91297 Arpajon Cedex France — We present ab initio evaluations of the thermal and of the electrical conductivity of hydrogen at a density of 80 g/cc [1] and 160 g/cc corresponding to the Inertial Confinement Fusion regime. Results are compared with different theories (Hubbard, Lee-More, Ichimaru) and with an average atom (AA) model coupled with a Kubo-Greenwood evaluation of transport coefficients. The Lorentz number, which is the ratio between the thermal to the electrical conductivity, given by the Wiedemann-Frantz law is checked in different regimes ranging from the highly degenerate to the kinetic one. We have also used ab initio calculations to compute the CH thermal conductivity and compared it to the AA approach using mixing rules.

[1] V. Recoules, F. Lambert, A. Decoster, B. Canaud and J. Clerouin, “Ab initio determination of thermal conductivity of dense hydrogen plasmas,” Phys. Rev. Letters 102, 075002 (2009).

Jean Clerouin
CEA/DAM/DIF 91297 Arpajon Cedex France

Date submitted: 20 Jul 2009

Electronic form version 1.4