

Abstract Submitted
for the DPP09 Meeting of
The American Physical Society

Study of fast electron transport in warm dense plasma targets¹

M.S. WEI, H. SAWADA, S. CHAWLA, N. NAKANII, D.P. HIGGINSON, B.S. PARADKAR, T. YABUUCHI, F.N. BEG, UC San Diego, K.U. AKLI, R.B. STEPHENS, General Atomics, A. MACPHEE, D. HEY, S. LE PAPE, H.S. MCLEAN, M. FOORD, M.K. KEY, P.K. PATEL, A.J. MACKINNON, LLNL, H. FRIESEN, H. TIEDJE, Y.Y. TSUI, R. FEDOSEJEVS, Univ. of Alberta, Canada, J. PASLEY, Univ. of York, UK, A. MORACE, D. BATANI, Univ. of Milano, Italy — Recent experiments [Le Pape et al., RSI 79, 106104, 2008] successfully demonstrated creation of a warm dense plasma using ns laser driven shock compression in a low-density foam target. Fast electron transport in such shock heated targets relevant to fast ignition is investigated using the Titan laser at LLNL. K-shell x-ray emission from the Cu fluorescent layer in the foam package targets is measured using Bragg crystal imagers and HOPG spectrometers. Electron transport through cold, conductor and shocked compressed foam targets is compared. The experiment is modeled using a 2D radiation hydro code and the hybrid PIC code LSP.

¹Work supported by US DOE under contracts DE-FC02-04ER54789 (Fusion Science Center) and DE-FG02-05ER54834 (ACE).

M.S. Wei
UC San Diego

Date submitted: 17 Jul 2009

Electronic form version 1.4