

Abstract Submitted
for the DPP09 Meeting of
The American Physical Society

Particle Control and Carbon Transport Experiments on DIII-D¹

S.L. ALLEN, LLNL, E.A. UNTERBERG, ORISE, N.H. BROOKS, M.A. MAHDAVI, GA, D.G. WHITE, MIT, J.W. DAVIS, A.A. HAASZ, B. FITZPATRICK, U. Toronto — As part of the 2009 Joint Research Target, DIII-D, along with MIT and NSTX have completed a series of experiments on particle control and transport. On DIII-D, both dynamic (calculated particle balance) and static (pumps closed or regenerated) particle balance experiments were carried out in both L- and H-mode with cryopumping. We find that the exhaust obtained from both techniques is comparable. We find that the uptake can be large in ohmic and L-mode, but in H-mode, the wall retention flux is very small, which is promising for long pulse burning plasma experiments. Also, in support of tritium control research for ITER, we have started a process to qualify the DIII-D internal components for an anticipated air bake (350°, 10 Torr) in 2010. We find issues with some copper components, but most components are compatible. Air baking removes the co-deposited carbon that can be rich in tritium in ITER with carbon walls.

¹Work supported by the US DOE under DE-AC52-07NA27344, DE-AC05-06OR23100, and DE-FC02-04ER54698.

Steve Allen
Lawrence Livermore National Laboratory

Date submitted: 20 Jul 2009

Electronic form version 1.4