

Abstract Submitted
for the DPP09 Meeting of
The American Physical Society

Progress on the Collisional-Radiative Modeling and the Analysis of Local Line Emission Profiles of Carbon Impurities from TJ-II Plasmas¹

R. FLORIDO, D. SUAREZ, R. RODRIGUEZ, J.M. GIL, J.G. RUBIANO, P. MARTEL, Departamento de Fisica, Universidad de Las Palmas de Gran Canaria, Spain, E. MINGUEZ, Instituto de Fusion Nuclear, Universidad Politecnica de Madrid, Spain, B. ZURRO, A. BACIERO, Laboratorio Nacional de Fusion, Asociacion Euratom-CIEMAT, Madrid, Spain — In a simple previous analysis the ratio of local emission profiles of carbon impurities (C^{4+} at 227.1 nm and C^{5+} at 529.0 nm) measured from TJ-II plasmas was found to have a significant dependence of neutral density profile. Hence, the aim of this work is to have a better theoretical understanding of this system and finally validate or discard the carbon spectral line ratio as a useful tool to estimate the neutral density profile in ECRH TJ-II plasmas. To assist in the theoretical modeling of this carbon system we use the collisional-radiative code ABAKO, which has been successfully applied to different ICF problems. Progress on the collisional-radiative modeling and simulation of MCF TJ-II plasmas are reported, paying particular attention to the influence of the charge-exchange recombination processes on the charge state distribution and carbon line ratio radial profiles.

¹Work supported by Research Project ENE2008-06668-C02/FTN of the Spanish Ministry of Science and Innovation.

R. Florido
Depto de Fisica, Universidad de Las Palmas de Gran Canaria,
35017 Las Palmas de Gran Canaria, Spain

Date submitted: 21 Jul 2009

Electronic form version 1.4