

Abstract Submitted  
for the DPP11 Meeting of  
The American Physical Society

**Rotation in the Plasma Flow Frame of Marginal Neoclassical Tearing Mode Islands in DIII-D and NSTX<sup>1</sup>** R.J. LA HAYE, R.J. BUTTERY, General Atomics, S.P. GERHARDT, PPPL, S.A. SABBAGH, Columbia U. — Small island effects inhibit the pervasive occurrence of neoclassical tearing modes.  $m/n = 2/1$  or  $3/1$  islands are reduced (by decreasing beta and thus the destabilizing bootstrap current density) to the self-stabilization size (“marginal point”) in DIII-D and NSTX [1]. Non-zero island propagation in the plasma flow frame can produce a stabilizing polarization current provided it occurs in the direction of (but does not exceed) the ion diamagnetic drift [2]. This “polarization threshold” naturally scales as a characteristic island size of several times the ion banana width, which is consistent with experiment. Comparison of measurements in DIII-D and NSTX discharges of similar cross-section (but different aspect ratio) with polarization current theory will be presented.

[1] R.J. La Haye, et al., Proc. 38th EPS Conf. on Plasma Phys., Strasbourg, France, 2011, P2.088.

[2] K. Imada and H.R. Wilson, Proc. 38th EPS Conf. on Plasma Phys., Strasbourg, France, 2011, O3.116.

<sup>1</sup>Work supported in part by US DOE under DE-FC02-04ER54698, DE-AC02-09CH11466, and DE-FG02-04ER54761.

Rob La Haye  
General Atomics

Date submitted: 28 Jul 2011

Electronic form version 1.4