

Abstract Submitted
for the DPP12 Meeting of
The American Physical Society

Extended gyrokinetic field theory for time-dependent magnetic confinement fields¹ HIDEO SUGAMA, TOMOHIKO WATANABE, MASANORI NUNAMI, National Institute for Fusion Science — In conventional gyrokinetic theories, the gyrocenter phase-space variables are defined by using the background magnetic confinement field that is assumed to be independent of time. Recently, several studies have been trying to perform long-time gyrokinetic turbulent transport simulations up to the transport time scale although they still use the above-mentioned assumption. However, the background or equilibrium magnetic field changes along with the pressure profile on the transport time scale. Therefore, in order to accurately describe the long-time behaviors of the gyrokinetic turbulence, we need to treat the time-dependent background field and show how to determine its time dependence. In this work, the gyrokinetic field theory [1] is extended to derive the condition which determines the time-dependent magnetic confinement fields in axisymmetric toroidal systems. Then, conservation laws of energy and momentum including the part of the time-dependent background fields are naturally derived from the extended gyrokinetic field theory.

[1] H. Sugama, Phys. Plasmas **7**, 405 (2000).

¹This work is supported in part by the Japanese Ministry of Education, Culture, Sports, Science and Technology (Grant Nos. 21560861 and 22760660).

Hideo Sugama
National Institute for Fusion Science

Date submitted: 10 Jul 2012

Electronic form version 1.4