

Abstract Submitted
for the DPP12 Meeting of
The American Physical Society

The Dynamics of Turbulence and Flow During the L-H Transition¹ Z. YAN, G.R. MCKEE, University of Wisconsin-Madison, J.A. BOEDO, D.L. RUDAKOV, G.R. TYNAN, P.H. DIAMOND, University of California San Diego, R.J. GROEBNER, T.H. OSBORNE, General Atomics, G. WANG, L. SCHMITZ, University of California Los Angeles — Comprehensive 2D turbulence and flow measurements are obtained before, during and after the L-H transition. An ion gyro-radius and density scan was performed on DIII-D to investigate the physics of the L-H transition and the threshold dependence on B_T and n_e . The amplitude of long wavelength density fluctuations scale approximately with ρ^* . Stronger poloidal turbulence flow shear is found at low density. The poloidal turbulence velocity spectrum changes from GAM at ~ 300 ms before the L-H transition to zonal flows near the transition. About $100 \mu\text{s}$ before the transition, the Reynolds stress gradient, inferred from 2D BES velocimetry, increases rapidly. The poloidal velocity and shear likewise peak near the same time, consistent with the prediction that Reynolds stress drives a zonal flow that triggers the L-H. The turbulence and shear dynamics leading to a typical L-H transition will also be compared with limit-cycle-oscillations between L- and H-modes observed under certain discharge conditions.

¹Work supported by the US Department of Energy under DE-FG02-08ER54999, DE-FG02-89ER53296, DE-FG02-07ER54917, DE-FC02-04ER54698 and DE-FG03-08ER54984.

Zheng Yan
University of Wisconsin-Madison

Date submitted: 13 Jul 2012

Electronic form version 1.4