

Abstract Submitted  
for the DPP12 Meeting of  
The American Physical Society

**Laboratory Experiments of Electromagnetic Velocity Shear-driven Instabilities**<sup>1</sup> ERIK TEJERO, WILLIAM AMATUCCI, CHRISTOPHER CRABTREE, GURUDAS GANGULI, Naval Research Laboratory, CHRISTOPHER COTHRAN, Sotera Defense Solutions — *In situ* observations of sheared plasma flows collocated with electromagnetic wave activity have led to a laboratory effort to investigate the impact of electromagnetic, velocity shear-driven instabilities on the near-Earth space plasma dynamics. Results from laboratory experiments will be presented that demonstrate strongly localized DC electric fields perpendicular to the ambient magnetic field can behave as a radiation source for electromagnetic ion cyclotron waves, transporting energy away from the region of wave generation. The transition from electrostatic to electromagnetic ion cyclotron (EMIC) wave propagation has been investigated under scaled ionospheric conditions. The general wave characteristics and wave dispersion experimentally observed are in agreement with theory. In addition, the electromagnetic component of these waves increased by two orders of magnitude as the plasma  $\beta$  was increased. The observed EMIC waves are predominantly azimuthally propagating  $m=1$  cylindrical waves, which propagate in the direction of the  $\mathbf{E} \times \mathbf{B}$  drift. Experimental observations and comparison to theory will be presented.

<sup>1</sup>This work is supported by the Naval Research Laboratory Base Program and NASA.

Erik Tejero  
Naval Research Laboratory

Date submitted: 13 Jul 2012

Electronic form version 1.4