Abstract Submitted for the DPP13 Meeting of The American Physical Society

Progress on Bayesian Inference of the Fast Ion Distribution Function¹ L. STAGNER, W.W. HEIDBRINK, X. CHEN, UC-Irvine, W. SALEWSKI, Tech. U. Denmark, B.A. GRIERSON, PPPL — The fast-ion distribution function (DF) has a complicated dependence on several phase-space variables. The standard analysis procedure in energetic particle research is to compute the DF theoretically, use that DF in forward modeling to predict diagnostic signals, then compare with measured data. However, when theory and experiment disagree (for one or more diagnostics), it is unclear how to proceed. Bayesian statistics provides a framework to infer the DF, quantify errors, and reconcile discrepant diagnostic measurements. Diagnostic errors and weight functions that describe the phase space sensitivity of the measurements are incorporated into Bayesian likelihood probabilities. Prior probabilities describe physical constraints. This poster will show reconstructions of classically described, low-power, MHD-quiescent distribution functions from actual FIDA measurements. A description of the full weight functions will also be shown.

¹This work is supported in part by the US Department of Energy under SC-G903402, DE-FC02-04ER54698 and DE-AC02-09CH11466

L. Stagner University of California-Irvine

Date submitted: 12 Jul 2013 Electronic form version 1.4