

Abstract Submitted  
for the DPP13 Meeting of  
The American Physical Society

### Cyclotron Mode Frequency Shifts in Multi-Species Ion Plasmas<sup>1</sup>

M. AFFOLTER, F. ANDEREGG, D.H.E. DUBIN, C.F. DRISCOLL, UCSD — We observe cyclotron modes varying as  $\cos(m_\theta\theta - \omega_m t)$  on laser-diagnosed multi-species ion plasma columns. These mode frequencies are shifted away from the “bare” cyclotron frequency  $\Omega_s = (q_s/M_s)B$  for each species  $s$ , due to electric fields and plasma rotation. For radially uniform ion species, we observe resonances at frequencies  $\omega_s = \Omega_s + [m_\theta - 2 + \delta_s(1 - R_m)]\omega_{ExB}$ , in close agreement with theory [1]. Here,  $\omega_{ExB}$  is the measured ExB rotation frequency,  $\delta_s$  is the relative fraction of the resonant species, and  $R_m$  is a small image charge correction. These frequency shifts are measured on the  $m_\theta = 1$  center-of-mass mode, on the  $m_\theta = 2$  “elliptical” mode, and on the novel  $m_\theta = 0$  radial “breathing” mode. Absent laser diagnostics, these mode frequencies could be used to estimate the plasma composition  $\delta_s$  and rotation  $\omega_{ExB}$ . When the plasma is laser cooled to  $10^{-5} < T < 10^{-2}$  eV, the ions centrifugally separate, each species concentrates toward “full density” over a narrow radial annulus, and all frequencies approach the  $\delta_s = 1$  limit. We also note that at high temperatures, i.e.  $T > 10^{-1}$  eV, the cyclotron resonances develop a multiple peak structure with temperature-dependent frequency spacing, and a connection to radially standing Bernstein modes [1] will be pursued.

[1] D.H.E. Dubin, Phys. Plasmas **20**, 042120 (2013).

<sup>1</sup>Supported by PHY-0903877, DE-SC0002451, DE-SC0008693.

Matthew Affolter  
UCSD

Date submitted: 12 Jul 2013

Electronic form version 1.4