

Abstract Submitted  
for the DPP13 Meeting of  
The American Physical Society

**Examination of the Evolution of Edge Profiles and Ion Transport Between ELMs on DIII-D**<sup>1</sup> J.-P. FLOYD, W.M. STACEY, S.C. MEL-LARD, Georgia Tech, R.J. GROEBNER, General Atomics — The measured edge profile evolution between edge-localized modes (ELMs) of DIII-D H-mode discharge 144977 is analyzed with a fine time resolution in order to understand the evolution of diffusive (inferred thermal diffusivities and ion diffusion coefficients) and non-diffusive (particle pinch velocities) transport properties in the edge, using the momentum balance methodology of Ref. [1]. The evolution of the density, temperature, rotation, and radial electric field profiles in the edge pedestal between ELMs provides information of these diffusive and non-diffusive transport processes in the pedestal of H-mode plasmas. This methodology is incorporated into the GTEDGE code developed for DIII-D data interpretation. Using a smaller integration time for the charge-exchange-recombination measurements than past analyses allows a more detailed examination of the time evolution of the ion temperature and rotation velocities.

[1] W.M. Stacey and R.J. Groebner, Nucl. Fusion **51**, 063024 (2011).

<sup>1</sup>Work supported by the US DOE under DE-FG01-ER54538 and DE-FC02-04ER54698.

J.-P. Floyd  
Georgia Tech

Date submitted: 12 Jul 2013

Electronic form version 1.4